



## METABOLISABLE ENERGY

### Testing at Hill Laboratories

#### Introduction

“Metabolisable Energy” (ME) is a concept used to characterise the nutritional value of animal feedstuffs. It is an estimate of the energy available to an animal from digestion of a feed material, expressed in units of megaJoules per kilogram of feed (MJ/kg DM). Feed materials are comprised of a number of chemical components (fats, proteins, carbohydrates etc), each of which are potentially metabolisable by an animal and contribute to the energy content of the feed. These components do not all contribute equally (for instance fats have higher energy content than proteins) and the concept of ME has been developed to account for these differences.




Figure 1. The total energy value of feed consumed by the animal is partitioned into waste and metabolisable energy by the digestion process, and then partitioned into energy for maintenance and for growth/production by metabolic processes.

Testing for ME has become important to New Zealand farmers as a means of comparing the nutritive value of animal feed materials. It is considered that ME is a better indicator of a feed's ability to support animal production than the simple measure of dry matter content.

#### The measurement of ME

ME can only be measured directly using carefully designed animal feeding trials. In these experiments, measurements of feed intake, wastes excreted, liveweight changes, and heat evolved are combined to calculate the total energy produced through digestion and metabolism of the feed.

Research has shown a strong relationship between the *in-vivo* digestibility of a feed (i.e. the proportion of the feed assimilated by the animal) and its ME value, and conversion equations have been derived relating the two. Feed digestibility can also be measured in the laboratory using synthetic enzymes to emulate the digestion process within the animal. This *in-vitro* digestibility can be converted to an *in-vivo* digestibility by calibration against standard materials used in animal feeding trials.

#### Hamilton

1 Clyde Street  
Hamilton 3216  
Private Bag 3205  
Hamilton 3240  
New Zealand  
T +64 7 858 2000  
F +64 7 858 2001

#### Christchurch

101c Waterloo Road  
Hornby Christchurch 8042  
PO Box 16607  
Christchurch 8441  
T +64 3 377 7176  
F +64 3 377 7276

There are some limitations with this approach.

- The laboratory *in-vitro* digestibility measurement mimics the feed digestion that might be achieved by animals, but it cannot take account of factors such as palatability, animal species and condition, dietary balance, etc.
- The standard materials used to convert *in-vitro* digestibility results to *in-vivo* values are a source of inconsistency in results, with some laboratories still using in-house proprietary standards.
- Components in some feed materials may not be effectively digested by synthetic enzymes, leading to an under-estimation of feed digestibility (and hence ME). These materials can include feed concentrates such as high fat content materials, palm kernel extract, fishmeal, and compound feeds.

## The measurement of digestibility/ME at Hill Laboratories

Hill Laboratories has adopted the procedure developed by the Australian Fodder Industry Association (AFIA), which uses a pepsin-cellulase digestion method and a set of readily available standard materials for the conversion of digestibility result from *in-vitro* to *in-vivo* values. The AFIA procedure provides standardisation and consistency among Australasian feed testing laboratories.

The AFIA digestibility test requires 5 days to complete, and NIR technology provides a rapid alternative, provided the feed sample is similar to those present in the NIR calibration set.

## Typical Feed Values

The following table gives general information on the feed quality of a range of typical forages.

**Table 1 Feed Quality for Forage Samples**

Feed Type	Dry Matter (%)	Crude Protein (%)	Acid Det. Fibre (%)	Neutral Det.Fibre (%)	Digestibility (%DOMD)	Metabolisable Energy (MJ/kg)
Mixed Pasture	12 – 25	20 – 30	20 – 30	30 – 45	65 – 80	9 – 12
Pasture silage	25 – 30	14 – 20	20 – 35	30 – 45	65 – 75	9 – 11
Cereal Silage	35 – 40	8 – 12	25 – 40	35 – 60	55 – 65	9 – 10.5
Maize Silage	25 – 35	6 – 9	25 – 35	35 – 50	60 – 70	9.5 – 11
Lucerne Foliage	15 – 25	20 – 30	25 – 30	35 – 45	60 – 70	9 – 12
Lucerne Hay	85 – 90	18 – 25	25 – 35	35 – 45	55 – 65	8 – 11

## Summary

- ME is a concept, rather than a directly measurable chemical component.
- ME may be superior to “Dry Matter” as an indicator of a feed’s value in animal maintenance and production.
- ME can only be measured using animal feeding trials, but may be estimated with an enzyme “digestibility” procedure.
- ME results from different laboratories may vary.
- Hill Laboratories uses the digestibility method recommended by the AFIA.
- NIR is used to give rapid estimates of AFIA digestibility and in turn ME.

## Further reading

1. AFRC (1993) Energy and Protein Requirements of Ruminants. An advisory manual prepared by the AFRC Technical Committee on Responses to Nutrients. CAB INTERNATIONAL, Wallingford, UK.
2. NRC Nutrient Requirements of Dairy Cattle. National Academy Press, Washington DC. 7th Revised Edition, 2001
3. Givens, D.I et al (editors) Forage Evaluation in Ruminant Nutrition. CABI, Wallingford, UK, 2000 Holmes, C.W., Brookes, I.M., Garrick, D.J., MacKenzie, D.D.S., Parkinson, T.J., Wilson, G.F. Milk Production from Pasture – Principles and Practices. Massey University, NZ, 2002

## Contact Details

For further information about any of the above tests please contact our client service managers.

Fiona Calvert	Agriculture Division Client Services Manager	fiona.calvert@hill-labs.co.nz
Janice Christiansen	Agriculture Division Client Services Manager	janice.christiansen@hill-labs.co.nz
Nevan Ofsoski	Agriculture Division Client Services Manager	nevan.ofsoski@hill-labs.co.nz
Jane Smillie	Agriculture Division Client Services Manager	jane.smillie@hill-labs.co.nz