



R J Hill Laboratories Ltd  
1 Clyde St, Private Bag 3205, HAMILTON  
tel: 07 8582000  
fax: 07 8582001  
email: S&P@hill-labs.co.nz  
internet: www.hill-labs.co.nz



## Sampling

Asparagus leaf samples can be collected just as the spears are emerging, or later when they are in fern. The latter growth stage is the preferred sampling period.

### Fern

**Sampling Time:** February to March.


**Plant Part:** Fern.

**Collect From:** Upper 30 cm portion of the fern.

**Quantity per Sample:** 10 - 15 ferns.

**Recommended Tests:** Basic Plant (BP).

**Comments:** To help diagnose an obvious problem, leaves showing the first signs of the distinctive symptoms should be collected as soon as abnormalities appear. If sampling outside the normal sampling time it is useful to take a second sample of similar, healthy leaves from nearby unaffected plants for analysis as a comparative standard.



### Spear

**Sampling Time:** At spear emergence.


**Plant Part:** Spear.

**Collect From:** Upper 9 cm of the spear tips.

**Quantity per Sample:** 20 - 30 spears.

**Recommended Tests:** Basic Plant (BP).

**Comments:** This is not the recommended sampling time, but samples can be taken if a problem is suspected.



### Soil

**Sampling Time:** Prior to crop establishment and annually at any time of the year, although autumn to early winter is recommended.

**Core Depth:** 15 cm.

**Collect From:** Throughout the asparagus beds.

**Quantity per Sample:** 12 - 20 cores.

**Recommended Tests:** Basic Soil (BS).

**Comments:** When sampling prior to orchard establishment, a 15 - 40 cm depth sample should also be taken, primarily to check the sub-soil pH.

If trying to diagnose a problem with crop growth and yield, samples should be collected from the rooting zones of the worst affected plants. In these circumstances, a second sample taken for comparative purposes from the rooting zones of healthy plants may be useful.

## Interpretation

Interpretation of the laboratory's results is possible by comparison with normal levels expected for the crop in question. The interpretation given here are based on the best information available and relate specifically to the sampling instructions given.

Fern			Spear			Soil		
Element	Unit	Normal Range	Element	Unit	Normal Range	Element	Unit	Normal Range
Nitrogen	(%)	2.4 - 3.8	Nitrogen	(%)	6.5 - 9.0	pH	-	6.0 - 6.7
Phosphorus	(%)	0.21 - 0.35	Phosphorus	(%)	0.60 - 0.90	Olsen P	(ug/ml)	30 - 50
Potassium	(%)	1.5 - 2.4	Potassium	(%)	3.0 - 4.6	Potassium	(me/100g)	0.60 - 1.00
Sulphur	(%)	0.25 - 0.40	Sulphur	(%)	0.70 - 0.90	Calcium	(me/100g)	6.0 - 12.0
Calcium	(%)	0.40 - 0.80	Calcium	(%)	0.30 - 0.40	Magnesium	(me/100g)	1.00 - 3.00
Magnesium	(%)	0.15 - 0.20	Magnesium	(%)	0.15 - 0.20	Sodium	(me/100g)	0.00 - 0.50
Sodium	(%)	0.02 - 0.10	Sodium	(%)	0.01 - 0.05	CEC	(me/100g)	12.0 - 25.0
Iron	(ug/g)	40 - 150	Iron	(ug/g)	50 - 150	Volume Weight	(g/ml)	0.60 - 1.00
Manganese	(ug/g)	25 - 100	Manganese	(ug/g)	20 - 40			
Zinc	(ug/g)	10 - 60	Zinc	(ug/g)	80 - 120			
Copper	(ug/g)	6 - 12	Copper	(ug/g)	7 - 20			
Boron	(ug/g)	50 - 100	Boron	(ug/g)	50 - 100			

### Comments:

Asparagus is a lime loving crop, preferring a soil pH above 6.0.

Asparagus is considered to have low phosphorus requirement, compared to other vegetable crops. It is, however, considered susceptible to boron deficiency.

Results for copper, zinc and manganese in leaves sprayed with fungicides will not be reliable due to adhering spray residues on the leaves.

Iron deficiency symptoms may exist even when leaf levels appear satisfactory. This may be due to the presence of physiologically inactive forms of iron within the tissue. Also, soil contamination of leaves growing near the ground may elevate total iron results.

### References

- Weir, R.G. and Cresswell, G.C. 1995. Plant nutrient disorders 3. Vegetable crops. Inkata Press.
- Reuter, D. J. and Robinson, J. B. (Eds) 1997. Plant analysis. An interpretation manual. Second edition.
- Blackmore, L.C; Searle, P.L and Daly, B.K. 1987. Methods for chemical analysis of soils. NZ Soil Bureau Scientific Report 80. NZ Soil Bureau, DSIR.
- Fertiliser recommendation for horticultural crops. HortResearch HortNET, 1997.
- Annual report of ministry of agriculture and fisheries. 1977, pg 177.
- Geraldson, C.M.; Klacan, G.R. and Lorenz, O.A. 1973. Soil testing and plant analysis. L. Walsh (Ed), Soil Science Society of America, Ch 22.

### Disclaimer:

Normal Range levels quotes relate specifically to the sampling procedure given. The Normal Range levels and Comments provided are the most up to date levels available but may be altered without notification. Such alterations are implemented immediately in the laboratory histogram reports. It is recommended that a consultant or crop specialist be involved with interpretations and recommendations.