



CROP GUIDE - Choumoellier


Sampling Notes

KB Item: 3450v3

The nutritional status of this vegetable crop is monitored using soil tests and plant analysis. Monitoring regularly is important to help sustain optimum levels and avoid nutritional disorders. If disorders do occur, rapid diagnosis is necessary to assist correction.

Leaf

Sampling Time:	Mid growth.
Plant Part	Youngest mature leaf.
Collect From:	-
Quantity per Sample:	20 leaves.
Recommended Tests:	Basic Plant (BP), Molybdenum (Mo).



Comments: This guide is to establish whether any nutrient imbalances exist to limit plant growth. An alternative approach is to evaluate its feed value for grazing livestock. In this case, the samples collected should be typical of the animals intake and include some stem material as well.

To help diagnose an obvious problem, leaves showing the first signs of the distinctive symptoms should be collected as soon as abnormalities appear. If sampling outside the normal sampling time it is useful to take a second sample of similar, healthy leaves from nearby unaffected plants for analysis as a comparative standard.

Soil

Sampling Time:	Prior to crop establishment
Core Depth	15cm
Collect From:	Randomly throughout the area to be planted
Quantity per Sample:	12 - 20 cores
Recommended Tests:	Basic Soil (BS), Sulphate Sulphur (SO ₄), Available Nitrogen (AN)

Comments: If a problem is suspected during the growing season, then a sample should be taken from the rooting zone immediately adjacent to the plant. Collecting a second sample from an unaffected area may help identify the cause of the problem.

Comments

Sulphur and molybdenum deficiencies are known to occur in brassica crops.

Nitrogen deficiency will appear as a purple pink coloration in the foliage. This symptom can also be induced by other factors such as cold weather, root damage from nematodes, drought stress and water logging.

Iron deficiency symptoms may exist even when leaf levels appear satisfactory. This may be due to the presence of physiologically inactive forms of iron within the tissue. Also, soil contamination of leaves growing near the ground may elevate total iron results.

References

Blackmore, L.C; Searle, P.L and Daly, B.K. 1987. Methods for chemical analysis of soils. NZ Soil Bureau Scientific Report 80. NZ Soil Bureau, DSIR.

Disclaimer

Normal Range levels shown as histograms in test reports relate specifically to the sampling procedure provided in this crop guide. The Normal Range levels in test reports and Comments provided in this Crop Guide are the most up to date available, but may be altered without notification. Such alterations are implemented immediately in the laboratory histogram reports. It is recommended that a consultant or crop specialist be involved with interpretations and recommendations.