



CROP GUIDE - Blackcurrant

Sampling Notes


KB Item: 3434v4

The nutritional status of blackcurrants is monitored using soil tests and plant analysis. Regular monitoring is important to help sustain optimum levels and avoid nutritional disorders. If disorders do occur, rapid diagnosis is necessary to assist correction.

Leaf

- Sampling Time:** During the 2 - 3 weeks prior to fruit ripening.
Plant Part: Youngest mature leaf (blade & petiole).
Collect From: Mid portion of the current season's extension growth.
Quantity per Sample: 5 representative leaves from each of 10 vines.
Recommended Tests: Basic Plant (BP).

Comments: To help diagnose an obvious problem, leaves showing the first signs of the distinctive symptoms should be collected as soon as abnormalities appear. If sampling outside the normal sampling time it is useful to take a second sample of similar, healthy leaves from nearby unaffected plants for analysis as a comparative standard.



Soil

- Sampling Time:** Prior to crop establishment and annually at any time of the year, although autumn to early winter is recommended.
Core Depth: 15cm.
Collect From: From the rooting zone of the plants.
Quantity per Sample: 12 - 20 cores.
Recommended Tests: Basic Soil (BS), Sulphur profile (S), Available Nitrogen (AN).
Comments: Soil samples are usually collected for analysis prior to planting the crop.

If trickle irrigation is used, the wetted zones of the soil should be sampled separately, as minerals in the water may produce abnormal test levels.

If trying to diagnose a problem with crop growth and yield, samples should be collected from the rooting zones of the worst affected plants. In these circumstances, a second sample taken for comparative purposes from the rooting zones of normal plants may be useful.



Comments

Most nutritional work on blackcurrants has been done in Britain and Eastern Europe, with little local information being available.

Foliar deficiency symptoms are rare in New Zealand, the most likely one being for nitrogen. Symptoms are bright orange, red or purplish red leaves.

Potassium deficiency symptoms start as red purplish leaves gradually going to necrotic marginal scorching.

Calcium and magnesium requirements are low.

Blackcurrants are sensitive to high levels of chloride, so chloride forms of fertiliser should be minimised.

Crop performance is best with a soil pH above 5.8.

References

Doughty, C. Horticulturist, Washington State University. Pers Comm.

Ballinger, W.E. and Goldstone, E.F. 1967. North Carolina, Agric. Expt. Tech. Bulletin, No. 178.

Fertiliser recommendation for horticultural crops. HortResearch HortNET, 1997.

Blackmore, L.C; Searle, P.L and Daly, B.K. 1987. Methods for chemical analysis of soils. NZ Soil Bureau Scientific Report 80. NZ Soil Bureau, DSIR.

Reuter, D. J. and Robinson, J. B. (Eds) 1997. Plant analysis. An interpretation manual. Second edition.

Mills, H. A. and Jones, Jr, J B 1996. Plant analysis handbook II.

Disclaimer

Normal Range levels shown as histograms in test reports relate specifically to the sampling procedure provided in this crop guide. The Normal Range levels in test reports and Comments provided in this Crop Guide are the most up to date available, but may be altered without notification. Such alterations are implemented immediately in the laboratory histogram reports. It is recommended that a consultant or crop specialist be involved with interpretations and recommendations.