

Triticale

Sampling Notes

Triticale is a hybrid between rye and wheat. Plant growth stage has a major influence on the nutrient levels in the tissue. Two distinct growth stages are specified for sample collection; neither preferred over the other, though each is useful as detailed below.

Triticale is quite similar to wheat except it has spreading growth until stem elongation, when the stems extend in the normal erect growth form of wheat.

Leaf (1) Late Tiller (GS25-GS29)

Sampling Time: When the leaves have formed, and the leaf-sheaths are lengthening and becoming erect. Just prior to extension.

Plant Part: Whole above portion of the plant.

Collect From: Random sites throughout the sampling area.

Quantity per Sample: 30 to 40 plants.

Recommended Tests: Basic Plant (BP).

Comments: This is the recommended plant sampling stage. The advantage of sampling at this early stage is that there may be time to correct nutrient disorders observed in the current crop.



Leaf (2) Ear Emergence (GS51-GS59)

Sampling Time: When stem extension is complete and the head of the ear emerges from the boot.

Plant Part: Top 30cm of the plant.

Collect From: Random sites throughout the sampling area.

Quantity per Sample: 20 to 30 plants.

Recommended Tests: Basic Plant (BP).

Comments: Testing at this later stage will indicate more accurately that the crop has accumulated the required nutrients successfully.

Soil

Sampling Time: Prior to crop establishment.

Core Depth: 15cm.

Collect From: Random sites throughout the sampling area.

Quantity per Sample: 12-20 cores.

Recommended Tests: Basic Soil (BS), Sulphur profile (S), Available Nitrogen (AN).

Comments: Soil samples are usually collected for analysis prior to planting the crop.

If trying to diagnose a problem with crop growth and yield, samples should be collected from the rooting zones of the worst affected areas. In these circumstances, a second sample taken for comparative purposes from the rooting zones of normal areas may be useful.

Comments

Most cultural techniques for growing wheat can be transferred directly to triticale.

Small grain production and quality are greatly influenced by fertilisation.

Nitrogen has been found to be the most important fertiliser element in New Zealand cereal crops. Significant responses to potassium, sulphur or magnesium have also been recorded.

Triticale is yield responsive and well adapted to high fertility conditions, such as effluent paddocks.

Improper growth stage identification can result in errors in interpretation. Nutrient uptake precedes dry matter accumulation occurring between tillering and ear emergence. Consequently, nutrient concentrations generally decline between these stages.

Diagnosis of sulphur deficiency can be assisted by using the N:S ratio. A sulphur deficiency may exist when the N:S ratio is greater than 16:1. Severe deficiency is likely when the ratio is greater than 20:1.

References

Jones Jr, J.B 1967. Soil testing and plant analysis. Part 2. SSSA Special Publication Series, p 49-58.

Ward, R.C.; Whitney, D.A. and Westfall, D.G. 1973. Plant analysis as an aid in fertilising small grains. Soil testing and plant analysis.

Lockman, R.B. 1969. Agronomy Abstracts, American Society of Agronomy, Wisconsin, pg 97.

Blackmore, L.C; Searle, P.L and Daly, B.K. 1987. Methods for chemical analysis of soils. NZ Soil Bureau Scientific Report 80. NZ Soil Bureau, DSIR.

Reuter, D. J. and Robinson, J. B. (Eds) 1997. Plant analysis. An interpretation manual. Second edition.

D. Salmon, M. McLelland, T. Schoff, P. Juskiw. 2001. Triticale Production Manual. Alberta Agriculture, Food and Rural Development (AAFRD) Agdex 118/20-1.

Disclaimer

Normal Range levels shown as histograms in test reports relate specifically to the sampling procedure provided in this crop guide. The Normal Range levels in test reports and Comments provided in this Crop Guide are the most up to date available, but may be altered without notification. Such alterations are implemented immediately in the laboratory histogram reports. It is recommended that a consultant or crop specialist be involved with interpretations and recommendations.
